Visions:

The Coming Revolutions in Particle Physics

Chris Quigg
Fermilab

Third Symposium on LHC Physics and Detectors

Chia, Sardinia (Italy) · October 27, 2001

Eppur si muove ...

Eppur si muove ...

> Completing the Copernican Revolution

Humans do not occupy a privileged location in the Universe

Eppur si muove ...

Completing the Copernican Revolution
 Humans do not occupy a privileged location in the Universe

Cimenti ...

Eppur si muove ...

Completing the Copernican Revolution
 Humans do not occupy a privileged location in the Universe

Cimenti ...

Learning to read Nature by doing experiments

Eppur si muove ...

Completing the Copernican Revolution
 Humans do not occupy a privileged location in the Universe

Cimenti ...

Rejecting Authority:Learning to read Nature by doing experiments

The minute particular ...

Eppur si muove ...

Completing the Copernican Revolution
 Humans do not occupy a privileged location in the Universe

Cimenti ...

▷ Rejecting Authority:Learning to read Nature by doing experiments

The minute particular ...

Not asking general questions and receiving limited answers, but asking limited questions and finding general answers

Our Picture of Matter

Pointlike $(r \lesssim 10^{-18} \text{ m})$ quarks

$$\left(\begin{array}{c} u \\ d \end{array} \right)_L \quad \left(\begin{array}{c} c \\ s \end{array} \right)_L \quad \left(\begin{array}{c} t \\ b \end{array} \right)_L$$

and **leptons** (idealization that neutrinos are massless) ...


$$\left(\begin{array}{c} \nu_e \\ e^- \end{array} \right)_L \quad \left(\begin{array}{c} \nu_\mu \\ \mu^- \end{array} \right)_L \quad \left(\begin{array}{c} \nu_ au \\ au^- \end{array} \right)_L$$

with interactions specified by

$${f SU(3)_c}\otimes {f SU(2)_L}\otimes {f U(1)_Y}$$

gauge symmetries ...

QCD explains the light hadron masses



For p, ρ , $[\pi]$, ..., confinement energy is the source. "Mass without mass"

From the 1898–99 University of Chicago catalogue:

"While it is never safe to affirm that the future of the Physical Sciences has no marvels in store even more astonishing than those of the past, it seems probable that most of the grand underlying principles have been firmly established and that further advances are to be sought chiefly in the rigorous application of these principles to all the phenomena which come under our notice An eminent physicist has remarked that the future truths of Physical Science are to be looked for in the sixth place of decimals."

Electroweak theory has many successes


- Neutral currents
- Charm
- Weak gauge bosons W^{\pm} and Z^0

A decade of LEP, et al.

- Testing the quantum field theory at the one per mille level
- Looking for new physics "in the sixth place of the decimals"

Precision measurements to determine unknown parameters . . .

Inferring the top-quark mass through its rôle in quantum corrections:



Precision measurements test the theory ...





The Result


NuTeV Measures:

$$\sin^2 \theta_W^{\text{(on-shell)}} = 0.2277 \pm 0.0013(\text{stat.}) \pm 0.0009(\text{syst.})$$

$$- 0.00022 \times (\frac{M_{\text{top}}^2 - (175 \text{ GeV})^2}{(50 \text{ GeV})^2})$$

$$+ 0.00032 \times \ln(\frac{M_{\text{Higgs}}}{150 \text{ GeV}})$$

cf. standard model fit (LEPEWWG), 0.2227 ± 0.00037 A discrepancy of $3\sigma...$



 $R_{\rm exp}^{\nu}$ and $R_{\rm exp}^{\overline{\nu}}$ measured to a precision of $0.3\%,\,0.65\%,\,{\rm respectively}$ (systematics lead to correlated uncertainty)


NuTeV's Technique

Charm Production and Charm Sea Errors are Large

⇒ Need a Technique Insensitive to Sea Quarks


Paschos-Wolfenstein Relation:

$$R^{-} = \frac{\sigma_{NC}^{\nu} - \sigma_{NC}^{\overline{\nu}}}{\sigma_{CC}^{\nu} - \sigma_{CC}^{\overline{\nu}}}$$
$$= \rho^{2} \left(\frac{1}{2} - \sin^{2}\theta_{W}\right)$$



- ullet R^- manifestly insensitive to sea quarks
 - \hookrightarrow Massive quark production enters from d_V quarks only (Cabbibo suppressed and at high x)
 - → Charm, strange sea errors negligible
- Requires Separate ν and $\overline{\nu}$ Beams \Rightarrow NuTeV SSQT


NuTeV Beamline



- Beam is almost purely ν or $\overline{\nu}$: $(\overline{\nu} \text{ in } \nu \text{ mode } 3 \times 10^{-4}, \ \nu \text{ in } \overline{\nu} \text{ mode } 4 \times 10^{-3})$
- Beam is $\sim 1.6\%$ electron neutrinos

uN Deep Inelastic Scattering at NuTeV

Lab E Detector - Fermilab E815(NuTeV)




Target/Calorimeter:

- 168 Fe plates (3m \times 3m \times 5.1cm)
- 84 liquid scintillation counters
 Trigger the detector
 Visible energy
 Neutrino interaction point
 Event length
- 42 drift chambers
 Localized transverse shower position

Toroidal Spectrometer:

• 15kG field $(P_T = 2.4 \, GeV/c)$



Scintillator


2" Steel

Continuous Test Beam: interspersed with ν beam

Hadron, muon and electron beams
 Map toroid and calorimeter response

SM Fit with NuTeV $\sin^2 \theta_W$

Fall 2001




(Courtesy M. Grunewald, LEPEWWG)

Without NuTeV: $\chi^2/\mathrm{dof} = 21.5/14$, probability of 9.0% With NuTeV: $\chi^2/\mathrm{dof} = 30.5/15$, probability of 1.0% Upper m_{Higgs} limit weakens slightly

Comparison with M_W


$$\sin^2 \theta_W^{\text{(on-shell)}} \equiv 1 - \frac{M_W^2}{M_Z^2}$$




- ullet In standard electroweak theory, NuTeV precision is comparable to a single direct measurement of M_W
- ullet More inconsistent with direct M_W than other data


Neutral Current ν Interactions

- ullet LEP I measures Z lineshape and decay partial widths to infer the "number of neutrinos"
 - \hookrightarrow Their result is $N_{\nu}=3\frac{\Gamma_{\exp}(Z\to \nu\overline{\nu})}{\Gamma_{\rm SM}(Z\to \nu\overline{\nu})}=3\times(0.9947\pm0.0028)$
 - \hookrightarrow LEP I "direct" partial width $(\nu\nu\gamma) \Rightarrow N_{\nu} = 3 \times (1.00 \pm 0.02)$
- $\stackrel{(-)}{\nu}_{\mu} e^- \rightarrow \stackrel{(-)}{\nu}_{\mu} e^-$ scattering (CHARM II *et al.*)
 - ightharpoonup PDG fit: $g_V^2 + g_A^2 = 0.259 \pm 0.014$, cf. 0.258 predicted
- NuTeV can fit for a deviation in $\nu\&\overline{\nu}$ NC rate
 - $\rightarrow \rho_0^2 = 0.9884 \pm 0.0026 \text{(stat)} \pm 0.0032 \text{(syst)}$




- In this interpretation, NuTeV confirms and strengthens LEP I indications of "weaker" neutrino neutral current
 - → NB: This is not a unique or model-independent interpretation!





Analogy to superconductivity sets M_W, M_Z



Meissner effect: EM fields disturb condensate of Cooper pairs Weak bosons disturb Higgs condensate, acquire masses:

$$M_W^2 = \frac{g^2 v^2}{2} = \frac{\pi \alpha}{G_F \sqrt{2} \sin^2 \theta_W}$$

$$M_Z^2 = \frac{M_W^2}{\cos^2 \theta_W}$$

EW scale is
$$v = (G_F\sqrt{2})^{-\frac{1}{2}} \approx 246 \text{ GeV}$$

Disturbing EW condensate may generate fermion mass

EWSB is necessary, not sufficient

Standard model: each fermion mass \Rightarrow new, unknown Yukawa coupling

$$\mathcal{L}_{\text{Yukawa}}^{(e)} = -\zeta_e \left[\overline{\mathsf{R}}(\phi^{\dagger} \mathsf{L}) + (\overline{\mathsf{L}}\phi) \mathsf{R} \right].$$

$$m_e = \zeta_e v / \sqrt{2}$$

All fermion masses \sim physics beyond the standard model!

$$\zeta_t \approx 1$$
 $\zeta_e \approx 3 \times 10^{-6}$ $\zeta_\nu \approx 10^{-10}$??

What accounts for the range and values of the Yukawa couplings?

There may be other sources of neutrino mass

Best hope until now:

Unified theories: pattern of fermion masses simplifies on high scales

YuKawa couplings offer the possibility of C.P violation...

CONSIDER

THE CP VIOLATION WITHIN THE SM (ARISING FROM THE QUARK MIXING MATRIX/YUKAWA COUPLINGS) IS ITSELF A WINDOW ON PHYSICS BEYOND THE STANDARD MODEL.



ν_{solar} News

$$\phi_{\text{SNO}}^{\text{CC}}(\nu_e) = 1.75 \pm 0.07 \text{ (stat.)}_{-0.11}^{+0.12} \text{ (sys.)} \pm 0.05 \text{ (th.)} \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$


$$\phi_{\text{SNO}}^{\text{ES}}(\nu_x) = 2.39 \pm 0.0.34 \text{ (stat.)}_{-0.14}^{+0.16} \text{ (sys.)} \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

$$\phi_{\text{Super-K}}^{\text{ES}}(\nu_x) = 2.32 \pm 0.03 \text{ (stat.)}_{-0.07}^{+0.08} \text{ (sys.)} \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$


$$\phi_{\text{Super-K}}^{\text{ES}}(\nu_x) - \phi_{\text{SNO}}^{\text{CC}}(\nu_e) = 0.57 \pm 0.17 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

 $\Longrightarrow \overline{\nu_{\mu}} \text{ and } \overline{\nu_{\tau}} \text{ arrive at Earth (at } 3.3\sigma)$

Combined SNO + Super-K ⁸B Neutrinos



Quantum corrections suggest the Higgs-boson mass ...




... within the standard electroweak theory







A light Higgs boson hints new physics nearby



Elementarity

> Are quarks and leptons structureless?

Elementarity

> Are quarks and leptons structureless?

Symmetry

- ▷ Electroweak symmetry breaking and the 1-TeV scale
- ▷ Origin of gauge symmetries

Elementarity

> Are quarks and leptons structureless?

Symmetry

- ▷ Electroweak symmetry breaking and the 1-TeV scale
- ▷ Origin of gauge symmetries

Unity

- > Coupling constant unification
- □ Unification of quarks and leptons (new forces!); of constituents and force particles
- ▷ Incorporation of gravity

Elementarity

> Are quarks and leptons structureless?

Symmetry

- ▷ Electroweak symmetry breaking and the 1-TeV scale
- ▷ Origin of gauge symmetries

Unity

- □ Unification of quarks and leptons (new forces!); of constituents and force particles
- ▷ Incorporation of gravity

Identity

- > Fermion masses and mixings; CP violation; neutrino oscillations
- > What makes an electron an electron and a top quark a top quark?

Elementarity

> Are quarks and leptons structureless?

Symmetry

- ▷ Electroweak symmetry breaking and the 1-TeV scale
- > Origin of gauge symmetries

Unity

- > Coupling constant unification
- □ Unification of quarks and leptons (new forces!); of constituents and force particles
- > Incorporation of gravity

Identity

- > Fermion masses and mixings; CP violation; neutrino oscillations
- > What makes an electron an electron and a top quark a top quark?

Topography

What is the fabric of space and time? . . . the origin of space and time?

The Vacuum Energy Problem (Veltman)

Higgs potential $V(\psi^{\dagger}\psi) = \mu(\psi^{\dagger}\psi) + 1\lambda I(\psi^{\dagger}\psi)^2$

At the minimum

$$V(\langle \varphi^{\dagger} \varphi \rangle_{o}) = \frac{\mu^{2} v^{2}}{4} = -\frac{|\lambda| v^{4}}{4} < 0$$

IDENTIFY MH = -2 M2.

Higgs potential contributes a field-independent constant energy density

$$SH = \frac{M_H^2 v^2}{8} > 10^8 \text{ GeV}$$
 $M_H > 100 \text{ GeV}$

Observed Vacuum Energy Density


Svac < 10⁻⁴⁶ GeV⁴

MISMATCH BY 54 ORDERS OF MAG.

How to zero SH?

Natural to neglect gravity in particle physics

$$G_{
m Newton}$$
 small $\iff M_{
m Planck} = \left(\frac{\hbar c}{G_{
m Newton}} \right)^{\frac{1}{2}} pprox 1.22 imes 10^{19} \ {
m GeV}$ large



Estimate
$$B(K \to \pi G) \sim \left(\frac{M_K}{M_{\rm Planck}}\right)^2 \sim 10^{-38}$$

300 years after Newton: Why is gravity weak?

Resolve the hierarchy problem: EXTEND THE STANDARD MODEL

$$SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \left\{ egin{array}{ll} {
m composite Higgs boson} \\ {
m technicolor / topcolor} \\ {
m supersymmetry} \\ {
m } {
m ...} \end{array}
ight.$$

Resolve the hierarchy problem: EXTEND THE STANDARD MODEL

$$SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \left\{ egin{array}{ll} {
m composite Higgs boson} \\ {
m technicolor / topcolor} \\ {
m supersymmetry} \\ {
m } {
m ...} \end{array}
ight.$$

Novel speculation: change gravity to understand why $M_{\rm Planck}\gg v$

Resolve the hierarchy problem: EXTEND THE STANDARD MODEL

$$SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \begin{cases} \text{composite Higgs boson} \\ \text{technicolor / topcolor} \\ \text{supersymmetry} \\ \dots \end{cases}$$

Novel speculation: change gravity to understand why $M_{\rm Planck} \gg v$

 \triangleright String theory: extra dimensions assumed $R_{\rm ed} \simeq 1/M_{\rm Planck} \simeq 10^{-35}$ cm

Resolve the hierarchy problem: EXTEND THE STANDARD MODEL


$$SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \begin{cases} \text{composite Higgs boson} \\ \text{technicolor / topcolor} \\ \text{supersymmetry} \\ \dots \end{cases}$$


Novel speculation: change gravity to understand why $M_{\rm Planck} \gg v$

$$ightharpoonup$$
 String theory: extra dimensions assumed $R_{\rm ed} \simeq 1/M_{\rm Planck} \simeq 10^{-35}$ cm

$$\triangleright$$
 Gravity follows Newtonian force law down to $\lesssim 1~\text{mm}$

$$V(r) = -\int dr_1 \int dr_2 \frac{G_{\text{Newton}} \rho(r_1) \rho(r_2)}{r_{12}} \left[1 + \varepsilon_{\text{G}} \exp(-r_{12}/\lambda_{\text{G}}) \right]$$






1F GRAVITY PROPAGATES IN EXTRA DIMS dimensional analysis changes (Gauss)

> n extra dim., radius $R \Rightarrow$ $G_N \sim M_{Pl}^{-2} \sim M_s^{-n-2} R^{-n}$

If $M_s \sim 1$ TeV, $R \leq 1$ mm for $n \geq 2$ $M_{Pl} = M_s (M_s R)^{n/2}$


would mean that MPE results from a false extrapolation



Gravity's true scale could be near EW scale. (Subversive idea)



Might Extra Dimensions Explain the Range of Fermion Masses?



Arkani-Hamed, Schmaltz, and Mirabelli:

Different fermions ride different tracks in the $5^{\rm th}$ dimension Small offsets in the new coordinate \Rightarrow exponential differences in masses

The Great Lesson of Science in the Twentieth Century

The human scale is not privileged for understanding Nature . . .

and may even be disadvantaged

In a decade or two, we can hope to ...

Understand electroweak symmetry breaking Observe the Higgs boson Measure neutrino masses and mixings Establish Majorana neutrinos $(\beta \beta_{0\nu})$ Thoroughly explore CP violation in B decays Exploit rare decays (K, D, \dots) Observe neutron EDM, pursue electron EDM Use top as a tool Observe new phases of matter Understand hadron structure quantitatively Uncover the full implications of QCD Observe proton decay Understand the baryon excess Catalogue matter and energy of the universe Measure dark energy equation of state Search for new macroscopic forces Determine GUT symmetry

Detect neutrinos from the universe Learn how to quantize gravity Learn why empty space is nearly weightless Test the inflation hypothesis Understand discrete symmetry violation Resolve the hierarchy problem Discover new gauge forces Directly detect dark-matter particles Explore extra spatial dimensions Understand the origin of large-scale structure Observe gravitational radiation Solve the strong CP problem Learn whether supersymmetry is TeV-scale Seek TeV-scale dynamical symmetry breaking Search for new strong dynamics Explain the highest-energy cosmic rays Formulate the problem of identity

In a decade or two, we can hope to ...

Understand electroweak symmetry breaking Observe the Higgs boson Measure neutrino masses and mixings Establish Majorana neutrinos $(\beta \beta_{0\nu})$ Thoroughly explore CP violation in B decays Exploit rare decays (K, D, \dots) Observe neutron EDM, pursue electron EDM Use top as a tool Observe new phases of matter Understand hadron structure quantitatively Uncover the full implications of QCD Observe proton decay Understand the baryon excess Catalogue matter and energy of the universe Measure dark energy equation of state Search for new macroscopic forces Determine GUT symmetry

Detect neutrinos from the universe Learn how to quantize gravity Learn why empty space is nearly weightless Test the inflation hypothesis Understand discrete symmetry violation Resolve the hierarchy problem Discover new gauge forces Directly detect dark-matter particles Explore extra spatial dimensions Understand the origin of large-scale structure Observe gravitational radiation Solve the strong CP problem Learn whether supersymmetry is TeV-scale Seek TeV-scale dynamical symmetry breaking Search for new strong dynamics Explain the highest-energy cosmic rays Formulate the problem of identity

...and to learn the right questions to ask

A Decade of Discovery Ahead

In the midst of a revolution in our conception of Nature, we deal with fundamental questions about our world, including

- Are the quarks and leptons elementary or composite?
- What are the symmetries of Nature, and how are they hidden from us?
- Are there new forms of matter, like the superpartners suggested by supersymmetry?
- Are there more fundamental forces?
- What makes an electron an electron and a top quark a top quark?
- What is the dimensionality of spacetime?

Nothing is too wonderful to be true,

if it be consistent with the laws of nature . . .

Experiment is the best test

Michael Faraday



Fig. 2. Montage der letzten Sprühschutzkörper. Durchmesser 80 cm.

[T]here has been one transforming change over this thousand years. It is the adoption of the scientific method: the commitment to experiment, to test every hypothesis. But it is broader than science. It is the open mind, the willingness in all aspects of life to consider possibilities other than the received truth. It is openness to reason.

—Anthony Lewis, New York *Times*, 12/31/99

On the first emulsions from the Pic du Midi:

"It was as if, suddenly, we had broken into a walled orchard, where protected trees had flourished and all kinds of exotic fruits had ripened in great profusion."

—Cecil Frank Powell